

Pathological Response and Survival in Triple-Negative Breast Cancer Following Neoadjuvant Carboplatin plus Docetaxel



Priyanka Sharma¹, Sara López-Tarruella², José Angel García-Saenz³, Qamar J. Khan¹, Henry L. Gómez⁴, Aleix Prat^{5,6}, Fernando Moreno⁵, Yolanda Jerez-Gilarranz², Agustí Barnadas⁷, Antoni C. Picornell², María del Monte-Millán², Milagros González-Rivera⁸, Tatiana Massarrah², Beatriz Pelaez-Lorenzo⁹, María Isabel Palomero², Ricardo González del Val², Javier Cortés¹⁰, Hugo Fuentes-Rivera⁴, Denisse Bretel Morales⁴, Iván Márquez-Rodas², Charles M. Perou¹¹, Carolyn Lehn¹, Yen Y. Wang¹, Jennifer R. Klemp¹, Joshua V. Mammen¹, Jamie L. Wagner¹, Amanda L. Amin¹, Anne P. O'Dea¹, Jaimie Heldstab¹, Roy A. Jensen¹, Bruce F. Kimler¹, Andrew K. Godwin¹, and Miguel Martín²

Abstract

Purpose: Prognostic value of pathologic complete response (pCR) and extent of pathologic response attained with anthracycline-free platinum plus taxane neoadjuvant chemotherapy (NAC) in triple-negative breast cancer (TNBC) is unknown. We report recurrence-free survival (RFS) and overall survival (OS) according to degree of pathologic response in patients treated with carboplatin plus docetaxel NAC.

Patients and Methods: One-hundred and ninety patients with stage I–III TNBC were treated with neoadjuvant carboplatin (AUC6) plus docetaxel (75 mg/m²) every 21 days × 6 cycles. pCR (no evidence of invasive tumor in breast and axilla) and Residual cancer burden (RCB) were evaluated. Patients were followed for recurrence and survival. Extent of pathologic response was associated with RFS and OS using the Kaplan–Meier method.

Results: Median age was 51 years, and 52% were node-positive. pCR and RCB I rates were 55% and 13%, respec-

tively. Five percent of pCR patients, 0% of RCB I patients, and 58% of RCB II/III patients received adjuvant anthracyclines. Three-year RFS and OS were 79% and 87%, respectively. Three-year RFS was 90% in patients with pCR and 66% in those without pCR [HR = 0.30; 95% confidence interval (CI), 0.14–0.62; *P* = 0.0001]. Three-year OS was 94% in patients with pCR and 79% in those without pCR (HR = 0.25; 95% CI, 0.10–0.63; *P* = 0.001). Patients with RCB I demonstrated 3-year RFS (93%) and OS (100%) similar to those with pCR. On multivariable analysis, higher tumor stage, node positivity, and RCB II/III were associated with worse RFS.

Conclusions: Neoadjuvant carboplatin plus docetaxel yields encouraging efficacy in TNBC. Patients achieving pCR or RCB I with this regimen demonstrate excellent 3-year RFS and OS without adjuvant anthracycline. *Clin Cancer Res*; 24(23); 5820–9. ©2018 AACR.

Introduction

Triple-negative breast cancer (TNBC), which is defined by the lack of expression of estrogen receptor (ER), progesterone receptor (PgR), and absence of ERBB2 (HER2) overexpression and/or gene

amplification, accounts for 15% of all the breast cancers in the United States. Compared with other breast cancer subtypes, TNBC is associated with inferior long-term outcomes (1–3). Adjuvant chemotherapy reduces the risk of distant recurrence and death in

¹Division of Medical Oncology, University of Kansas Medical Center, Westwood, Kansas. ²Department of Medical Oncology, Hospital General Universitario Gregorio Marañón, Instituto de Investigación Sanitaria Gregorio Marañón, Universidad Complutense, CIBERONC, GEICAM, Madrid, Spain. ³Department of Medical Oncology, Hospital Clínico San Carlos, Madrid, Spain. ⁴Department of Medical Oncology, Instituto Nacional de Enfermedades Neoplásicas, Lima, Perú. ⁵Department of Medical Oncology, Hospital Clinic of Barcelona, Barcelona, Spain. ⁶Translational Genomics and Targeted Therapeutics in Solid Tumors, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain. ⁷Department of Medical Oncology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain. ⁸Laboratory of Translational Oncology, Instituto de Investigación Sanitaria Gregorio Marañón (IISGM), Madrid, Spain. ⁹Department of Oncology, Hospital Clínico, Valladolid, Spain. ¹⁰Department of Oncology, Ramón y Cajal University Hospital, Madrid, Spain. ¹¹Vall d'Hebron Institute of Oncology (VHIO), Barcelona, Spain. ¹²Departments of Genetics and Pathology & Laboratory Medicine, Lineberger Comprehensive Cancer Center, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina.

Note: Supplementary data for this article are available at Clinical Cancer Research Online (<http://clincancerres.aacrjournals.org/>).

Prior presentation: This work was presented in part at the 52nd Annual Meeting of the American Society of Clinical Oncology, June 3–7, 2016, Chicago, IL.

ClinicalTrials.gov: NCT02302742, NCT01560663

Corresponding Authors: Priyanka Sharma, University of Kansas Medical Center, 2330 Shawnee Mission Parkway, Mail Stop 5003, Westwood, KS 66205. Phone: 1-913-588-6079; Fax: 1-913-588-4085; E-mail: psharma2@kumc.edu; and Miguel Martín, Medical Oncology Service, Hospital General Universitario Gregorio Marañón, Facultad de Medicina, Universidad Complutense de Madrid, Instituto de Investigación Sanitaria Gregorio Marañón (IISGM), Dr Esquerdo 46, 28007 Madrid, Spain. Phone: 91-586-80-00; E-mail: mmartin@geicam.org

doi: 10.1158/1078-0432.CCR-18-0585

©2018 American Association for Cancer Research.

Translational Relevance

Phenotypic and molecular similarities between sporadic and *BRCA* mutation-associated triple-negative breast cancer (TNBC) have prompted the exploration of DNA-damaging agents like platinum compounds in TNBC. We report survival outcomes with neoadjuvant carboplatin plus docetaxel, an anthracycline-free chemotherapy regimen, in TNBC. Robust pathologic complete response (pCR) and near-complete response (RCB I) rates were noted with this regimen. Patients who achieved pCR or RCB I had excellent 3-year recurrence-free rate and overall survival (OS) without adjuvant anthracyclines, suggesting that the composite of pCR + RCB I may accurately identify patients at low risk of recurrence who can avoid further adjuvant chemotherapy and associated toxicities. This platinum-taxane chemotherapy regimen rendered equally high pCR and survival rates in *BRCA*-associated TNBC as in *BRCA* wild type TNBC. The results of our study support further evaluation of platinum-based chemotherapy in both *BRCA*-associated and wild type TNBC as well as the role of extent of pathologic response in risk stratification of patients undergoing neoadjuvant chemotherapy.

patients with TNBC and is generally recommended for TNBC patients with stage I ($T > 1$ cm)–III disease (4–6). Even so, despite receiving standard anthracycline-taxane-based chemotherapy, a significant proportion (20%–40%) of patients with early-stage TNBC develop metastatic disease (7, 8). Improved therapeutic approaches are thus desired for TNBC. In the absence of available actionable molecular targets, modifications of traditional chemotherapy regimens (i.e., the addition of platinum agents) may be potential means of improving outcomes for this breast cancer subtype.

Platinum compounds generate the double-stranded DNA breaks that are preferentially repaired by the mechanism of homologous recombination. Phenotypic and molecular similarities between *BRCA*-associated and sporadic TNBC suggest that homologous recombination deficiency may be a shared alteration that can be targeted in a larger subset of TNBC (7, 9–13). Germline *BRCA1/2* mutations are the prototype molecular alterations that confer homologous recombination deficiency and sensitivity to DNA-damaging therapy. Only 15% to 20% of patients with TNBC harbor germline *BRCA* mutations; however, additional mechanisms such as promoter methylation, transcript instability/attenuation, or somatic/germline mutations in other homologous recombination pathway genes may compromise DNA repair machinery (14–17). It has recently been reported that comprehensive evaluation of factors beyond germline *BRCA* mutation reveals homologous recombination deficiency in 50% to 60% of TNBC, expanding the therapeutic potential of DNA-damaging agents like platinum compounds within the general population of TNBC (16, 18). Neoadjuvant studies demonstrate that the addition of carboplatin to anthracycline and taxane-based neoadjuvant chemotherapy (NAC) improves pathologic complete response (pCR) in TNBC (11, 19–22). However, the improvement in pCR rate comes at the cost of increased toxicity (11, 19, 21, 22). Exploration of a platinum-based chemotherapy regimen with an improved toxicity profile in TNBC is therefore warranted. Taxanes

demonstrate significant activity in TNBC as well as preclinical and clinical synergy with platinum agents, providing rationale for evaluation of a platinum-taxane combination in TNBC (23–28).

We have recently reported encouraging efficacy of a carboplatin plus docetaxel NAC regimen in TNBC (29). In this study, overall pCR rates were 55% and were not influenced by germline *BRCA* mutation status (pCR 59% in *BRCA*-associated TNBC, 56% in *BRCA* wild type TNBC). Here, we report 3-year recurrence-free survival (RFS) and overall survival (OS) from the same study population. We also present the correlation between survival outcomes and degree of pathologic response as assessed by residual cancer burden (RCB).

Patients and Methods

Patient population

The study population includes patients with stage I–III TNBC treated with a NAC regimen of carboplatin plus docetaxel in two separate and independent prospective cohorts. Details of the study population have been published previously (29). A brief description of the study population is provided below.

This study was conducted in accordance with the U.S. Common Rule and the International Ethical Guidelines for Biomedical Research Involving Human Subjects. Human Investigations were performed after approval by the Institutional Review Board at each institution, and all subjects provided written informed consent.

University of Kansas (KU) cohort. Patients with stage I ($T \geq 1$ cm), II, and III TNBC were enrolled in an IRB-approved multisite prospective registry protocol (NCT02302742). Triple negativity was defined as ER and PgR IHC nuclear staining of less than 10% and HER2 IHC staining of 0 to 1+ or FISH ratio < 2.0 if IHC 2+ or if IHC not performed (30). From 2011 to 2015, 69 enrolled patients with stage I ($T > 1$ cm), II, and III TNBC were treated with a NAC regimen of carboplatin plus docetaxel.

MMJ-CAR-2014-01 (Spanish) cohort. Female patients with pathologically confirmed diagnosis of stage II–III ($T \geq 2$ cm) TNBC were enrolled on a prospective, multicenter, nonrandomized trial exploring the antitumor activity of neoadjuvant carboplatin plus docetaxel (NCT01560663). See Supplementary Material for a list of participating institutions. Triple negativity was defined as ER and PgR nuclear staining of less than 1% by IHC and HER2 IHC staining of 0 to 1+ or FISH ratio < 2.0 if IHC 2+ or if IHC not performed (30). Between 2010 and 2015, 123 patients with stage II and III TNBC were enrolled and treated with neoadjuvant carboplatin plus docetaxel.

This study was conducted in accordance with the U.S. Common Rule and the International Ethical Guidelines for Biomedical Research Involving Human Subjects. Human investigations were performed after approval by the Institutional Review Board at each institution, and all subjects provided written informed consent.

Study procedures

As described previously (29), patients in both cohorts were prescribed an NAC regimen of carboplatin (AUC 6) + docetaxel (75 mg/m^2) given every 21 days for 6 cycles. All patients received myeloid growth factor support (KU cohort: 6 mg pegfilgrastim on D2, Spanish cohort: filgrastim $300 \text{ } \mu\text{g/day} \times 5$ to 7 days after chemotherapy, according to the guidelines of each institution). In

patients with clinically suspicious axillary lymph node/s, histologic confirmation by biopsy or fine-needle aspiration was encouraged. Following NAC, all patients underwent breast surgery. Axillary sampling was required except in patients with pretreatment negative sentinel lymph nodes. Twenty-two percent (41/183) of the per-protocol population underwent pretreatment sentinel lymph node biopsy; completion dissection was required in case of positive pretreatment sentinel lymph node biopsy (additional information regarding pretreatment axillary lymph node assessment is included as Supplementary Text). Extent of axillary surgery, subsequent irradiation, and postoperative adjuvant chemotherapy were otherwise determined by the treating physician. Patients were prospectively followed for recurrence and survival status.

Pathologic evaluation

As described previously (29), pathologic response was determined locally without central pathologic review. All surgical pathology reports were centrally reviewed by the principal investigators of the respective cohorts. pCR was defined as the absence of residual invasive disease in the breast and axilla with or without ductal carcinoma *in situ* (ypT0/isN0). Patients with pCR in the breast and negative pretreatment sentinel lymph nodes were considered to have achieved pCR. RCB for all the patients was calculated centrally (utilizing parameters reported on individual patient pathology reports), blinded to enrollment site, using the classification by Symmans *et al* (31). Patients achieving pCR (RCB 0) or near-pCR (RCB I) were assessed within the group RCB 0/I.

Germline BRCA1/2 testing

KU cohort. Germline testing for *BRCA1/2* was done utilizing commercially available tests and laboratories.

Spanish cohort. Germline testing for *BRCA1/2* was done at Sistemas Genómicos facilities using targeted next-generation sequencing of 7 genes (*BRCA1*, *BRCA2*, *PALB2*, *BARD1*, *RAD50*, *RAD51C*, and *RAD51D*) and multiplex ligation-dependent probe amplification (MLPA) by quantification of probes corresponding to *BRCA1* and *BRCA2* genes using the MLPA Kit (MRC-Holland) according to the manufacturer's recommendations, fragment analysis by ABI 3730xl genetic analyzer (Applied Biosystems), and data normal-

ization and interpretation of results using Coffalyser.net software as recommended by MRC-Holland. Only *BRCA1* and *BRCA2* results are available and reported for this analysis.

Patients with germline *BRCA1* or *BRCA2* mutation were counseled in both cohorts as per institutional guidelines.

Statistical analysis

Relevant demographic, treatment, and outcome variables of the Spanish and KU cohorts were combined for analysis. Demographic and toxicity data depict the intent-to-treat population, whereas all other analyses reflect the per-protocol patient population. RFS was defined as the time from diagnosis to first recurrence (invasive ipsilateral breast, invasive local/regional, or distant) or to death as a result of any cause (32). OS was defined as time from diagnosis to death as a result of any cause. Patients were censored on the date of last contact if an event had not been observed. Confidence intervals (CI) for the proportion of patients with pCR and RCB 0/I were calculated according to the exact two-sided binomial test. Survival curves were assessed by the Kaplan-Meier method, and unadjusted survival comparisons were conducted using log-rank tests. Cox regression modeling was used for univariate and multivariable analysis of factors associated with risk of recurrence and death. All reported *P* values and CIs are from two-sided tests. All analyses were conducted using SPSS Statistics version 22 (IBM Corporation).

Results

Study cohort

A CONSORT diagram depicting cohort identification has been provided (Fig. 1). As reported previously (29), the study population included 190 subjects treated in KU and Spanish cohorts between 2010 and 2015. Two patients were found to have metastatic disease after enrollment and were excluded from the intent-to-treat population. Seven patients in the intent-to-treat population were not included in the per-protocol analysis (consent withdrawn or lost to follow-up prior to definitive breast surgery).

Patient characteristics

Table 1 describes the demographic and baseline clinical characteristics of the study population. For the overall study




Figure 1. CONSORT diagram.^aIn addition to the various centers in Spain, the Spanish cohort includes a center from Lima, Peru.

Table 1. Patient characteristics

	All patients (N = 190)
Age at diagnosis, years - Median (range)	51 (29-81)
Race - n (%)	
Caucasian	174 (92%)
Black	14 (7%)
Asian	2 (1%)
Ethnicity ^a	
Hispanic	26 (14%)
Non-Hispanic	163 (86%)
Menopausal status	
Pre/peri	86 (45%)
Post	98 (52%)
Unknown	6 (3%)
Histologic grade	
1	1 (1%)
2	39 (21%)
3	140 (78%)
T stage	
1	23 (12%)
2	104 (55%)
3	34 (18%)
4	29 (15%)
Lymph node status	
Negative	90 (47%)
Positive	98 (52%)
Unknown	2 (1%)
ER/PR IHC	
0%	184 (97%)
1-10%	6 (3%)
Germline <i>BRCA</i> mutation	
Absent	136 (71%)
Present	30 (16%)
Unknown	24 (13%)
Surgery type ^b	
Lumpectomy	84 (45%)
Mastectomy	101 (55%)
pCR (n = 183) ^b	
Yes	100 (55%)
No	83 (45%)
RCB 0/I (n = 180) ^c	
Yes	123 (68%)
No	57 (31%)
Adjuvant chemotherapy in patients with RCB II+III (n = 57)	
Yes	33 (58%)
No	24 (42%)

^aEthnicity is not known for one patient.

^bSurgical information is not available for 5 patients.

^cRCB status is not available for 3 patients.

population, median age was 51 years (range, 29–81 years), 14% were Hispanic, 7% were black, and 52% had clinically node-positive disease. Sixteen percent of the study population carried a deleterious *BRCA1/2* mutation (germline *BRCA* testing results were available for 87% of patients). A small percentage (3%, all from KU cohort) had ER/PgR expression between 1% and 10%.

Pathologic response

Pathologic complete response and RCB I for the per-protocol study population (N = 183) were 55% (100/183; 95% CI, 48–62) and 13% (23/183; 95% CI, 8–18), respectively (Supplementary Fig. S1), as reported previously (29). Taken together, this yields a RCB 0/I rate of 68% (123/183; 95% CI, 61–75). When assessed by germline *BRCA* status, pCR rate was 56% (75/133) in *BRCA* wild type and 59% (16/27) in *BRCA* mutation-associated TNBC (P = 0.83). Because of the inclusion of a small number of patients with

1% to 10% ER/PgR expression, pCR assessment was also performed according to ER/PgR expression status. Fifty percent and 55% of patients with ER/PgR 1% to 10% (n = 6) and ER/PgR <1% (n = 177), respectively, achieved pCR (P = 1.0).

Survival outcomes


At a median follow-up of 36 months (range, 4–97 months), there have been 36 RFS events (distant n = 26, local/regional n = 5, site of metastasis unknown n = 5) and 25 deaths, with estimated 3-year RFS and OS of 79% (95% CI, 73–85) and 87% (95% CI: 82–92), respectively (Supplementary Fig. S2). Figure 2 shows Kaplan–Meier curves for RFS and OS by pCR (Fig. 2A and B) and RCB 0/I (Fig. 2C and D) status. Patients achieving pCR demonstrated significantly better RFS and OS compared with patients without pCR. Estimated 3-year RFS was 90% and 66% in patients with and without pCR, respectively (HR = 0.30; 95% CI, 0.14–0.62; P = 0.0001). Estimated 3-year OS was 94% (95% CI, 89–99) and 79% (95% CI, 70–88) for patients with and without pCR, respectively (HR = 0.25; 95% CI, 0.10–0.63; P = 0.001). Similar to those who achieved pCR, patients who achieved RCB 0/I status also demonstrated significantly superior RFS and OS compared with patients with RCB II/III. Estimated 3-year RFS was 91% (95% CI, 86–96) and 59% (95% CI, 45–73) in patients with RCB 0/I and RCB II/III, respectively (HR = 0.20; 95% CI: 0.10–0.41, P < 0.0001). Estimated 3-year OS was 95% (95% CI: 91–99) and 75% (95% CI: 63–87) for patients with RCB 0/I and RCB II/III, respectively (HR = 0.16; 95% CI: 0.06–0.42, P < 0.0001). Figure 3 shows RFS and OS by the four RCB classes. Estimated 3-year RFS was 90% (95% CI: 84–96) for RCB 0 (pCR), 93% (95% CI: 79–100) for RCB I, 69% (95% CI: 54%–84) for RCB II, and 21% (95% CI, 0–46) for RCB III. Estimated 3-year OS was 94% (95% CI, 89–99) for RCB 0 (pCR), 100% for RCB I, 83% (95% CI, 71–95) for RCB II, and 43% (95% CI, 11–75) for RCB III. Compared with patients with RCB II, patients with RCB III had inferior RFS (HR = 4.70; 95% CI, 1.97–11.20; P < 0.0001) and OS (HR = 4.34; 95% CI, 1.59–11.84; P = 0.002). RFS appeared similar between RCB 0 (pCR) and RCB I groups (HR = 2.48; 95% CI, 0.32–19.36; P = 0.37). Because of the absence of OS events in the RCB I group, no formal statistical analysis was done to compare OS between RCB 0 and RCB I groups.

When analyzed by *BRCA* status, estimated 3-year RFS was 88% (95% CI, 75–100) in patients with germline *BRCA1/2* mutation and 79% (95% CI, 72–86) in patients who were *BRCA* wild type (HR = 0.69; 95% CI, 0.24–1.99; P = 0.48). Similar to RFS, estimated 3-year OS was similar between patients with germline *BRCA1/2* mutation (95%; 95% CI, 86–100) and those who were *BRCA* wild type (87%, 95% CI, 81–93; HR = 0.37; 95% CI, 0.12–2.21; P = 0.33).

Adjuvant chemotherapy use and survival

Only 5 of 100 patients with pCR received postsurgical adjuvant chemotherapy (all 5 received doxorubicin and cyclophosphamide, AC). Adjuvant chemotherapy use in patients with pCR did not impact RFS or OS (data not shown). None of the 23 patients with RCB I status received adjuvant chemotherapy. Fifty-eight percent (33/57) of patients with RCB II/III received adjuvant anthracyclines, and the survival outcomes in this group were not significantly different compared with those patients who did not receive adjuvant anthracycline (Supplementary Fig. S3). No patients received adjuvant capecitabine.

Sharma et al.

**Figure 2.**

A, RFS by pCR status; **B**, OS by pCR status; **C**, RFS by RCB 0/I vs. RCB II/III; **D**, OS by RCB 0/I vs. RCB II/III. ^aRCB unavailable for 3 patients.

Factors impacting survival

On univariate analysis, baseline node-positive status, higher baseline T-stage, lack of pCR, and an RCB greater than 0/I were all predictors of a greater risk of recurrence and death, whereas age at diagnosis, *BRCA* mutation status, pathologic grade, and adjuvant chemotherapy use (in patients with RCB II/III) did not impact RFS and OS (Table 2). On multivariable analysis, higher baseline T-stage was associated with greater risk of recurrence and death, whereas RCB 0/I was associated with reduced risk of recurrence and death. Baseline node-positive status was associated with greater risk of recurrence but not of death.

Adverse events


As reported previously (29), 28% (54/190) of patients in the intent-to-treat population experienced one or more grade 3/4 adverse events (grade 3 = 21%, grade 4 = 7%; Table 3). Of the per-protocol population, 88% (161/183) of patients completed all six cycles of treatment. Twelve percent (22/183) discontinued treat-

ment prematurely: 4.4% (8/183) because of progressive disease, 6.0% (11/183) because of toxicity, 1.6% (3/183) because of other reasons. No treatment-related deaths were reported.

Discussion

In this study, we demonstrate that carboplatin plus docetaxel NAC regimen yields a high pathologic response rate (55%), which translates to encouraging 3-year RFS and OS. Recent randomized trials demonstrate that the addition of carboplatin to anthracycline and taxane-based NAC improves pathologic response from 37%–41% to 50%–58% in TNBC (11, 19, 21, 22, 27). The pCR rate of 55% with this carboplatin plus docetaxel regimen appears to be comparable with the pCR rate noted when carboplatin is added to anthracycline-taxane chemotherapy and numerically higher than classical neoadjuvant anthracycline-taxane combinations, where 28% to 40% of TNBC patients achieved pCR (11, 21, 33). Long-term benefits from the addition of neoadjuvant

Survival after Neoadjuvant Carboplatin + Docetaxel in TNBC

**Figure 3.**

A, RFS by RCB class; **B**, OS by RCB class. ^aRCB unavailable for 3 patients.

platinum to anthracycline/taxane regimens in TNBC are not yet clear. The Cancer and Leukemia Group B (CALGB) 40603 randomized phase II trial evaluated the addition of carboplatin and/or bevacizumab to a standard chemotherapy regimen of weekly paclitaxel followed by AC in TNBC and demonstrated improvement in pCR rates with the addition of carboplatin (41% vs. 54%; $P = 0.003$). In this trial, however, the addition of carboplatin did not lead to improvement in 3-year event-free survival (71% vs. 76%, $P = 0.36$; ref. 11). The GeparSixto randomized phase II trial reported that addition of weekly carboplatin to the combination of paclitaxel, nonpegylated liposomal doxorubicin, and bevacizumab improved pCR rate (37% vs. 53%, $P = 0.005$), and unlike CALGB 40603, at a median follow-up of 3 years demonstrated a 44% improvement in 3-year disease-free survival (DFS; 76% vs. 85%, $P = 0.0325$) in patients with TNBC (21, 34). It is possible that differences in the chemotherapy backbones of the control arm between CALGB 40603 and GeparSixto (i.e., absence of the alkylating agent cyclophosphamide in the control arm of GeparSixto) could have contributed to the differences in the long-term outcomes noted in the two studies. However, it is noteworthy that neither CALGB 40603 nor GeparSixto were sufficiently powered for DFS and OS endpoints; several ongoing, adequately powered, randomized phase III trials are evaluating the efficacy of adjuvant platinum in the context of AC-plus-taxane chemotherapy in TNBC (NCT02488967, NCT02455141, NCT02441933). In our study, where more than half of patients had node-positive disease and a third had stage III disease, we reported a 3-year RFS and OS of 79% and 87%, respectively. These findings are in line with 3-year outcomes reported from GeparSixto (48% node-positive) and CALGB 40603 (58% node-positive) and also compare favorably with outcomes noted in contemporary adjuvant trials of AC plus taxane in TNBC (35, 36).

In a recently reported germline *BRCA*-focused subgroup analysis of GeparSixto, pCR and DFS benefit from the addition of carboplatin was observed primarily in patients without germline *BRCA* mutation (37). *BRCA* mutation carriers experienced a high pCR rate of 66.7% with the anthracycline + taxane + bevacizumab regimen, and this rate was not increased further by addition

of carboplatin. In contrast, in patients with *BRCA* wild type TNBC, the addition of carboplatin increased pCR from 36% to 55%. Regardless of the treatment regimen, DFS was generally high in *BRCA* mutation carriers, with no significant difference separated by study arms, whereas in patients with *BRCA* wild type TNBC, better DFS was noted in the carboplatin arm (HR = 0.53; 95% CI, 0.29–0.96; $P = 0.04$). These findings suggest that the substantial homologous recombination deficiency induced by germline *BRCA* mutation may be therapeutically responsive to most types of DNA-damaging chemotherapy, including anthracyclines, and that this responsiveness is not improved further by addition of a platinum agent. However, the homologous recombination deficiency brought about by mechanisms other than germline *BRCA* mutation may not be as robust, and could therapeutically respond to targeting by platinum agents, even in the presence of anthracyclines. Further translational research is needed to understand mechanisms of homologous recombination deficiency in *BRCA* wild type TNBC and their relationship to platinum response.

Previous studies and meta-analyses have demonstrated that pCR is a robust surrogate for improved long-term outcomes in aggressive breast cancer subtypes like TNBC (3, 38). The positive impact of pCR on long-term outcomes (event/disease-free and OS) was also observed in both GeparSixto and CALGB 40603 trials. Hazard ratios for the prognostic impact of pCR on RFS and OS noted with the current carboplatin plus docetaxel regimen overlap with hazard ratios noted in the meta-analysis by Cortazar et al and those from both GeparSixto and CALGB 40603. It is noteworthy that for patients with pCR, where only 5% received adjuvant anthracycline-based chemotherapy, we demonstrate 3-year RFS and OS of 90% and 94%, respectively. These findings suggest that the favorable prognostic impact of pCR in TNBC is probably independent of the chemotherapy regimen leading to pCR, a finding which has also recently been noted by the I-SPY 2 investigators (39). An interesting finding of our trial is the excellent outcome of patients who achieved RCB class I pathologic response, which was as good as the outcome of those achieving pCR. A similar finding has recently been reported for TNBC patients treated with neoadjuvant anthracycline and taxane-based

Sharma et al.

Table 2. Univariate and multivariate analysis of RFS and OS

Variable	Univariate analysis			
	RFS		OS	
	HR (95% CI)	P	HR (95% CI)	P
Age				
≤50	1		1	
>50	0.90 (0.47-1.73)	0.75	0.60 (0.27-1.33)	0.20
Lymph node status				
Negative	1		1	
Positive	4.47 (1.96-10.21)	<0.0001	5.40 (1.85-15.74)	0.001
T stage				
1/2	1		1	
3/4	4.56 (2.30-9.02)	<0.0001	6.11 (2.55-14.65)	<0.0001
Histologic grade				
1/2	1		1	
3	1.18 (0.57-2.46)	0.66	0.56 (0.19-1.64)	0.26
Germline <i>BRCA</i> mutation				
Negative/unknown	1		1	
Positive	0.69 (0.24-2.00)	0.49	0.51 (0.12-2.21)	0.29
pCR				
No	1		1	
Yes	0.30 CI (0.14-0.61)	0.0005	0.25 (0.10-0.63)	0.001
RCB				
II/III	1		1	
0/I	0.20 (0.10-0.41)	<0.0001	0.16 (0.06-0.42)	<0.0001
RCB				
II/III	1		1	
I	0.09 (0.01-0.67)	0.003	0.027 (0.00-2.16)	0.006
Adjuvant chemotherapy (in patients with RCB II/III)				
Yes	1		1	
No	1.30 (0.56-3.03)	0.54	0.75 (0.27-2.08)	0.58
	Multivariate analysis ^a			
Variable	RFS		OS	
	HR (95% CI)	P	HR (95% CI)	P
RCB				
II/III	1		1	
0/I	0.30 (0.14-0.63)	0.002	0.26 (0.10-0.70)	0.008
T stage				
1/2	1		1	
3/4	2.28 (1.07-4.86)	0.032	2.79 (1.06-7.36)	0.038
Lymph node status				
Negative	1		1	
Positive	2.51 (1.04-6.03)	0.040	2.47 (0.79-7.74)	0.12

^aVariables included age, lymph node status, T stage, pathologic response, RCB class, germline mutation status, adjuvant chemotherapy.

NAC (40). If confirmed in additional studies, a composite of pCR plus RCB class I pathologic response could be more suitable than pCR alone for identification of TNBC patients destined for excellent prognosis following NAC. Two thirds of patients (68%) in this study demonstrated pCR or RCB I with this carboplatin plus

docetaxel regimen, and these patients enjoyed very robust 3-year RFS ($\geq 90\%$) and OS ($\geq 93\%$) without adjuvant anthracycline treatment. These data indicate that adjuvant anthracycline can be avoided for patients achieving pCR or RCB with carboplatin plus docetaxel.

Poor outcomes were noted in patients with RCB II and III, particularly so in patients with RCB III, where the 3-year RFS was only 21%. In our study, a modest proportion (58%) of patients with RCB II or III received adjuvant anthracycline-based chemotherapy; however, adjuvant chemotherapy use did not impact outcomes in these patients. These findings highlight the need for and support the ongoing investigations of novel therapies in TNBC patients who have significant residual disease.

Anthracyclines and cyclophosphamide, although very active for treatment of breast cancer, carry established small but serious long-term risks (secondary leukemia/myelodysplastic syndrome, cardiomyopathy; refs. 41, 42). Anthracycline-free platinum-taxane chemotherapy regimen is associated with lower incidence of secondary leukemia/myelodysplastic syndrome and cardiomyopathy compared with anthracyclines-cyclophosphamide-taxane regimen when given concurrent with trastuzumab in HER2-positive

Table 3. Grade 3 to 4 treatment-related toxicities

N = 190	Grade 3/4 n (%)	Grade 3 n (%)	Grade 4 n (%)
Leukopenia	2 (1%)	2 (1%)	0 (0%)
Neutropenia	22 (12%)	13 (7%)	9 (5%)
Thrombocytopenia	11 (6%)	9 (5%)	2 (1%)
Anemia	7 (4%)	6 (3%)	1 (1%)
Febrile neutropenia	8 (4%)	6 (3%)	2 (1%)
Nausea	5 (3%)	5 (3%)	0 (0%)
Vomiting	5 (3%)	4 (2%)	1 (1%)
Mucositis	2 (1%)	2 (1%)	0 (0%)
Diarrhea	5 (3%)	5 (3%)	0 (0%)
Peripheral neuropathy	3 (2%)	3 (2%)	0 (0%)
Fatigue	2 (1%)	2 (1%)	0 (0%)
Other ^a	9 (5%)	9 (5%)	0 (0%)

^aOther: Hepatic/transaminase elevation ($n = 3$), rash ($n = 2$), hyponatremia ($n = 1$), thrombosis ($n = 1$), allergic reaction ($n = 1$), dehydration ($n = 1$).

breast cancer (43). The efficacy of anthracycline-free neoadjuvant combinations in sporadic and *BRCA*-associated TNBC has not been adequately explored so far. Taxanes are important components of neo/adjuvant chemotherapy regimens for breast cancer treatment and appear to contribute particularly among patients with TNBC (23, 44). Platinum and taxanes are mechanistically distinct, and preclinical and clinical data demonstrate synergy between platinum compounds and taxanes in several solid tumor types, including metastatic TNBC (24–26, 28). In the metastatic setting, the randomized TNT trial demonstrated superior efficacy of single-agent carboplatin over single-agent docetaxel in the presence of germline *BRCA* mutation but equal efficacy of either agent in the absence of germline *BRCA* mutation. Furthermore, the TNT trial noted no cross-resistance between the two drugs, with equal response rates noted for both agents upon cross-over (45). The promising efficacy of the neoadjuvant carboplatin plus docetaxel regimen noted in the current study might also be explained by the notion that the combination of docetaxel and carboplatin offers robust antitumor coverage for both basal and nonbasal TNBC (45). The encouraging outcomes noted with this carboplatin plus docetaxel regimen provide further evidence for the clinical efficacy of platinum-taxane combination in TNBC. Together, these data provide clinical and biological rationale for further investigation of anthracycline-free, platinum-taxane chemotherapy for early-stage TNBC.

Tolerance of the carboplatin plus docetaxel regimen used in this study was good, with only 6% of patients discontinuing therapy because of toxicity. A similar chemotherapy regimen, the TCH regimen (that combines carboplatin and docetaxel with trastuzumab), is widely used for the neo/adjuvant treatment of HER2-positive breast cancer and is likewise associated with a favorable toxicity profile.

Sixteen percent of the current study population carried a deleterious germline *BRCA1/2* mutation, which is consistent with expected prevalence of germline *BRCA1/2* mutation in TNBC (14, 15). We noted a pCR rate of 59% in *BRCA* mutation carriers, which is in line with previously reported pCR rates with single-agent platinum chemotherapy in *BRCA1* mutation carriers (46). In our study, deleterious *BRCA* mutation was not associated with 3-year DFS or OS, a finding that is also consistent with previous reports (47, 48). Furthermore, it is becoming increasingly evident that 40% to 50% of *BRCA* wild type TNBC may harbor homologous recombination deficiency (as identified by genomic mutational and transcriptional signatures), and that the benefit of platinum agents may extend beyond *BRCA* mutation-associated breast cancers (12, 16, 17, 49, 50). Several homologous recombination deficiency assays are being evaluated in clinical and translational research studies; however, assays with demonstrable clinical utility in selecting *BRCA* wild type TNBC patients likely to derive preferential benefit from platinum agents are not yet available.

Our study has limitations. First, the decision to combine the two patient cohorts for the purpose of reporting pCR and survival outcomes was made *post hoc* and was not preplanned. However, this report includes almost 200 patients with TNBC treated uniformly across three different continents. In this combined analysis, more than half of the patients had node-positive disease, and a third had stage III disease. These patient characteristics are very similar to contemporary neo/adjuvant randomized clinical trials for TNBC (11, 36).

Second, we acknowledge the lack of a control arm, a deficit that can only be addressed in the setting of a randomized trial. In fact,

an ongoing randomized phase II study is currently comparing neoadjuvant carboplatin plus docetaxel regimen to an anthracycline-taxane-carboplatin regimen (NCT02413320). While we await data from ongoing trials, this regimen may provide an alternative for patients who desire treatment with a platinum regimen but are not candidates for anthracycline. We also acknowledge the modest follow-up of 3 years; however, given the natural history of TNBC, where the majority of recurrence events occur in the first 3 years, longer follow-up is unlikely to significantly change our results (2, 3).

In conclusion, the combination of carboplatin plus docetaxel yielded significant antitumor activity in stage I–III TNBC, with a favorable toxicity profile.

Disclosure of Potential Conflicts of Interest

F. Moreno is a consultant/advisory board member for Pfizer and Roche. J. Cortés is a consultant/advisory board member for AstraZeneca, Biothera, Celgene, Celleria Biotech, Eisai, Merus, Novartis, Pfizer, Roche, and Seattle Genetics. I. Márquez-Rodas is a consultant/advisory board member for Bristol-Myers Squibb, MSD, Novartis, Pierre Fabre, and Roche. C.M. Perou is an employee of and holds ownership interest (including patents) in Bioclassifier LLC. M. Martín is a consultant/advisory board member for AstraZeneca, Lilly, Novartis, Pfizer, and Roche. No potential conflicts of interest were disclosed by the other authors.

Authors' Contributions

Conception and design: P. Sharma, S. López-Tarruella, Q.J. Khan, C.M. Perou, J.R. Klemm, R.A. Jensen, M. Martín

Development of methodology: P. Sharma, S. López-Tarruella, Q.J. Khan, A. Prat, A.K. Godwin, M. Martín

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): P. Sharma, S. López-Tarruella, J.A. García-Saenz, Q.J. Khan, H. Gómez, A. Prat, F. Moreno, Y. Jerez-Gilarranz, A. Barnadas, M. González-Rivera, B. Pelaez-Lorenzo, M.I. Palomero, R.G. del Val, J. Cortes, H. Fuentes-Rivera, I. Márquez-Rodas, C. Lehn, Y.Y. Wang, J.R. Klemm, J.V. Mammen, J.L. Wagner, A.L. Amin, A.P. O'Dea, J. Heldstab, A.K. Godwin, M. Martín

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): P. Sharma, S. López-Tarruella, H. Gómez, A. Prat, A. Barnadas, A.C. Picornell, M. del Monte-Millán, I. Márquez-Rodas, C.M. Perou, Y.Y. Wang, J.R. Klemm, B.F. Kimler, A.K. Godwin, M. Martín

Writing, review, and/or revision of the manuscript: P. Sharma, S. López-Tarruella, J.A. García-Saenz, Q.J. Khan, A. Prat, F. Moreno, Y. Jerez-Gilarranz, A. Barnadas, T. Massarrah, M.I. Palomero, J. Cortés, H. Fuentes-Rivera, I. Márquez-Rodas, C.M. Perou, C. Lehn, J.R. Klemm, J.M. Mammen, A. Amin, A.P. O'Dea, R.A. Jensen, B.F. Kimler, A.K. Godwin, M. Martín

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): P. Sharma, D. Bretel Morales, Y.Y. Wang, R.A. Jensen, B.F. Kimler, A.K. Godwin, M. Martín

Study supervision: P. Sharma, S. López-Tarruella, M. Martín

Acknowledgments

This work was supported by the University of Kansas Research Career Award, University of Kansas Cancer Center's CCRG (P30 CA168524), and Biospecimen Repository Core Facility to P. Sharma; research grant from Instituto de Salud Carlos III (PI 12/02684) and funding from Centro de Investigación Biomédica en Red de Salud Carlos III to M. Martín; and National Cancer Institute Breast SPORE program (P50-CA58223) to C. M. Perou.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received February 19, 2018; revised May 21, 2018; accepted July 24, 2018; published first July 30, 2018.

References

- Carey LA, Dees EC, Sawyer L, Gatti L, Moore DT, Collichio F, et al. The triple negative paradox: primary tumor chemosensitivity of breast cancer subtypes. *Clin Cancer Res* 2007;13:2329–34.
- Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA, et al. Triple-negative breast cancer: clinical features and patterns of recurrence. *Clin Cancer Res* 2007;13:4429–34.
- Liedtke C, Mazouni C, Hess KR, Andre F, Tordai A, Mejia JA, et al. Response to neoadjuvant therapy and long-term survival in patients with triple-negative breast cancer. *J Clin Oncol* 2008;26:1275–81.
- Senkus E, Kyriakides S, Penault-Llorca F, Poortmans P, Thompson A, Zackrisson S, et al. Primary breast cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013;24:vi7–23.
- National Comprehensive Cancer Network. NCCN clinical practice guidelines in oncology: breast cancer. 2017. Available from: https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf.
- Coates AS, Winer EP, Goldhirsch A, Gelber RD, Gnant M, Piccart-Gebhart M, et al. Tailoring therapies-improving the management of early breast cancer: St Gallen International Expert Consensus on the primary therapy of early breast cancer 2015. *Ann Oncol* 2015;26:1533–46.
- Nielsen TO, Hsu FD, Jensen K, Cheang M, Karaca G, Hu Z, et al. Immunohistochemical and clinical characterization of the basal-like subtype of invasive breast carcinoma. *Clin Cancer Res* 2004;10:5367–74.
- Haffty BG, Yang Q, Reiss M, Kearney T, Higgins SA, Weidhaas J, et al. Locoregional relapse and distant metastasis in conservatively managed triple negative early-stage breast cancer. *J Clin Oncol* 2006;24:5652–7.
- Lips EH, Mulder L, Oonk A, van der Kolk LE, Hogervorst FB, Imholz AL, et al. Triple-negative breast cancer: BRCAness and concordance of clinical features with BRCA1-mutation carriers. *Br J Cancer* 2013;108:2172–7.
- Isakoff SJ, Mayer EL, He L, Traina TA, Carey LA, Krag KJ, et al. TBCRC009: a multicenter phase II clinical trial of platinum monotherapy with biomarker assessment in metastatic triple-negative breast cancer. *J Clin Oncol* 2015;33:1902–9.
- Sikov WM, Berry DA, Perou CM, Singh B, Cirincione CT, Tolane SM, et al. Impact of the addition of carboplatin and/or bevacizumab to neoadjuvant once-per-week paclitaxel followed by dose-dense doxorubicin and cyclophosphamide on pathologic complete response rates in stage II to III triple-negative breast cancer: CALGB 40603 (Alliance). *J Clin Oncol* 2015;33:13–21.
- Telli ML, Jensen KC, Vinayak S, Kurian AW, Lipson JA, Flaherty PJ, et al. Phase II study of gemcitabine, carboplatin, and iniparib as neoadjuvant therapy for triple-negative and BRCA1/2 mutation-associated breast cancer with assessment of a tumor-based measure of genomic instability: PrECOG 0105. *J Clin Oncol* 2015;33:1895–901.
- Prat A, Cruz C, Hoadley KA, Diez O, Perou CM, Balmana J. Molecular features of the basal-like breast cancer subtype based on BRCA1 mutation status. *Breast Cancer Res Treat* 2014;147:185–91.
- Sharma P, Klemp JR, Kimler BF, Mahnken JD, Geier LJ, Khan QJ, et al. Germline BRCA mutation evaluation in a prospective triple-negative breast cancer registry: implications for hereditary breast and/or ovarian cancer syndrome testing. *Breast Cancer Res Treat* 2014;145:707–14.
- Couch FJ, Hart SN, Sharma P, Toland AE, Wang X, Miron P, et al. Inherited mutations in 17 breast cancer susceptibility genes among a large triple-negative breast cancer cohort unselected for family history of breast cancer. *J Clin Oncol* 2015;33:304–11.
- Sharma P, Barlow W, Godwin A, Pathak H, Isakova K, Williams D, et al. Impact of homologous recombination deficiency biomarkers on outcomes in patients with triple-negative breast cancer treated with adjuvant doxorubicin and cyclophosphamide (SWOG S9313). *Ann Oncol* 2018;29:654–60.
- Mulligan JM, Hill LA, Deharo S, Irwin G, Boyle D, Keating KE, et al. Identification and validation of an anthracycline/cyclophosphamide-based chemotherapy response assay in breast cancer. *J Natl Cancer Inst* 2014;106:djt335.
- Telli ML, Timms KM, Reid J, Hennessy B, Mills GB, Jensen KC, et al. Homologous recombination deficiency (HRD) score predicts response to platinum-containing neoadjuvant chemotherapy in patients with triple-negative breast cancer. *Clin Cancer Res* 2016;22:3764–73.
- Rugo HS, Olopade OI, DeMichele A, Yau C, van't Veer LJ, Buxton MB, et al. Adaptive randomization of veliparib-carboplatin treatment in breast cancer. *N Engl J Med* 2016;375:23–34.
- Petrelli F, Coiu A, Borronovo K, Cabiddu M, Ghilardi M, Lonati V, et al. The value of platinum agents as neoadjuvant chemotherapy in triple-negative breast cancers: a systematic review and meta-analysis. *Breast Cancer Res Treat* 2014;144:223–32.
- von Minckwitz G, Schneeweiss A, Loibl S, Salat C, Denkert C, Rezaei M, et al. Neoadjuvant carboplatin in patients with triple-negative and HER2-positive early breast cancer (GeparSixto; GBG 66): a randomised phase 2 trial. *Lancet Oncol* 2014;15:747–56.
- Loibl S, O'Shaughnessy J, Untch M, Sikov WM, Rugo HS, McKee MD, et al. Addition of the PARP inhibitor veliparib plus carboplatin or carboplatin alone to standard neoadjuvant chemotherapy in triple-negative breast cancer (BrightNess): a randomised, phase 3 trial. *Lancet Oncol* 2018;19:497–509.
- Hayes DF, Thor AD, Dressler LG, Weaver D, Edgerton S, Cowan D, et al. HER2 and response to paclitaxel in node-positive breast cancer. *N Engl J Med* 2007;357:1496–506.
- McGuire WP, Hoskins WJ, Brady MF, Kucera PR, Partridge EE, Look KY, et al. Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer. *N Engl J Med* 1996;334:1–6.
- Engblom P, Rantanen V, Kulmala J, Helenius H, Grenman S. Additive and supra-additive cytotoxicity of cisplatin-taxane combinations in ovarian carcinoma cell lines. *Br J Cancer* 1999;79:286–92.
- Delbaldo C, Michiels S, Syz N, Soria JC, Le Chevalier T, Pignon JP. Benefits of adding a drug to a single-agent or a 2-agent chemotherapy regimen in advanced non-small-cell lung cancer: a meta-analysis. *JAMA* 2004;292:470–84.
- Gluz O, Nitz U, Liedtke C, Christgen M, Grischke EM, Forstbauer H, et al. Comparison of neoadjuvant nab-paclitaxel+carboplatin vs nab-paclitaxel+gemcitabine in triple-negative breast cancer: randomized WSG-ADAPT-TN trial results. *J Natl Cancer Inst* 2018;110:628–37.
- Yardley D, Coleman R, Conte P, Cortes J, Brufsky A, Shtivelband M, et al. Nab-paclitaxel + carboplatin or gemcitabine vs gemcitabine/carboplatin as first-line treatment for patients with triple-negative metastatic breast cancer: results from the randomized phase 2 portion of the nAcity trial [abstract]. In: Proceedings of the 2016 San Antonio Breast Cancer Symposium; 2016 Dec 6–10; San Antonio, TX. Philadelphia (PA): AACR. Abstract nr P5–15–03.
- Sharma P, Lopez-Tarruella S, Garcia-Saenz JA, Ward C, Connor CS, Gomez HL, et al. Efficacy of neoadjuvant carboplatin plus docetaxel in triple negative breast cancer: combined analysis of two cohorts. *Clin Cancer Res* 2017;23:649–57.
- Wolff AC, Hammond MEH, Hicks DG, Dowsett M, McShane LM, Allison KH, et al. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline update. *J Clin Oncol* 2013;31:3997–4013.
- Symmans WF, Peintinger F, Hatzis C, Rajan R, Kuerer H, Valero V, et al. Measurement of residual breast cancer burden to predict survival after neoadjuvant chemotherapy. *J Clin Oncol* 2007;25:4414–22.
- Hudis CA, Barlow WE, Costantino JP, Gray RJ, Pritchard KI, Chapman JA, et al. Proposal for standardized definitions for efficacy end points in adjuvant breast cancer trials: the STEEP system. *J Clin Oncol* 2007;25:2127–32.
- Arun B, Bayraktar S, Liu DD, Gutierrez Barrera AM, Atchley D, Pusztai L, et al. Response to neoadjuvant systemic therapy for breast cancer in BRCA mutation carriers and noncarriers: a single-institution experience. *J Clin Oncol* 2011;29:3739–46.
- von Minckwitz G, Loibl S, Schneeweiss A, Salat CT, Rezaei M, Zahm D-M, et al. Early survival analysis of the randomized phase II trial investigating the addition of carboplatin to neoadjuvant therapy for triple-negative and HER2-positive early breast cancer (GeparSixto) [abstract]. In: Proceedings of the Thirty-Eighth Annual CTRC-AACR San Antonio Breast Cancer Symposium; 2015 Dec 8–12; San Antonio, TX. Philadelphia (PA): AACR. Cancer Res 2016;76:Abstract nr S2–04.
- Sparano JA, Zhao F, Martino S, Ligibel JA, Perez EA, Saphner T, et al. Long-term follow-up of the E1199 phase III trial evaluating the role of taxane and schedule in operable breast cancer. *J Clin Oncol* 2015;33:2353–60.
- Yardley DA, Arrowsmith ER, Daniel BR, Eakle J, Brufsky A, Drosick DR, et al. TITAN: phase III study of doxorubicin/cyclophosphamide followed by

- ixabepilone or paclitaxel in early-stage triple-negative breast cancer. *Breast Cancer Res Treat* 2017;164:649–58.
37. Hahnen E, Lederer B, Hauke J, Loibl S, Krober S, Schneeweiss A, et al. Germline mutation status, pathological complete response, and disease-free survival in triple-negative breast cancer: secondary analysis of the GeparSixto randomized clinical trial. *JAMA Oncol* 2017;3:1378–85.
 38. Cortazar P, Geyer CE Jr. Pathological complete response in neoadjuvant treatment of breast cancer. *Ann Surg Oncol* 2015;22:1441–6.
 39. Yee D, DeMichele A, Isaacs C, Symmans F, yau C, Albain K, et al. Pathological complete response predicts event-free and distant disease-free survival in the I-SPY2 trial [abstract]. In: Proceedings of the 2017 San Antonio Breast Cancer Symposium; 2017 Dec 5–9; San Antonio, TX. Philadelphia (PA): AACR. Abstr GS3–08.
 40. Symmans WF, Wei C, Gould R, Yu X, Zhang Y, Liu M, et al. Long-term prognostic risk after neoadjuvant chemotherapy associated with residual cancer burden and breast cancer subtype. *J Clin Oncol* 2017;35:1049–60.
 41. Tan TC, Neilan TG, Francis S, Plana JC, Scherrer-Crosbie M. Anthracycline-induced cardiomyopathy in adults. *Compr Physiol* 2015;5:1517–40.
 42. Wolff AC, Blackford AL, Visvanathan K, Rugo HS, Moy B, Goldstein LJ, et al. Risk of marrow neoplasms after adjuvant breast cancer therapy: the national comprehensive cancer network experience. *J Clin Oncol* 2015;33:340–8.
 43. Slamon D, Eiermann W, Robert N, Giermek J, Martin M, Jasiowka M, et al. Ten year follow-up of BCIRG-006 comparing doxorubicin plus cyclophosphamide followed by docetaxel (AC→T) with doxorubicin plus cyclophosphamide followed by docetaxel and trastuzumab (AC→TH) with docetaxel, carboplatin and trastuzumab (TCH) in HER2+ early breast cancer [abstract]. *Cancer Res* 2016;76:Abstr S5–04.
 44. Jacquin JP, Jones S, Magne N, Chapelle C, Ellis P, Janni W, et al. Docetaxel-containing adjuvant chemotherapy in patients with early stage breast cancer. Consistency of effect independent of nodal and biomarker status: a meta-analysis of 14 randomized clinical trials. *Breast Cancer Res Treat* 2012;134:903–13.
 45. Tutt A, Tovey H, Cheang M, Kernaghan S, Kilburn L, Gazinska P, et al. Carboplatin in BRCA1/2-mutated and triple-negative breast cancer BRCAness subgroups: the TNT trial. *Nat Med* 2018;24:628–637.
 46. Byrski T, Huzarski T, Dent R, Marczyk E, Jasiowka M, Gronwald J, et al. Pathologic complete response to neoadjuvant cisplatin in BRCA1-positive breast cancer patients. *Breast Cancer Res Treat* 2014;147:401–5.
 47. Brekelmans CT, Tilanus-Linthorst MM, Seynaeve C, vd Ouweland A, Menke-Pluymers MB, Bartels CC, et al. Tumour characteristics, survival and prognostic factors of hereditary breast cancer from BRCA2-, BRCA1- and non-BRCA1/2 families as compared to sporadic breast cancer cases. *Eur J Cancer* 2007;43:867–76.
 48. Robson ME, Chappuis PO, Satagopan J, Wong N, Boyd J, Goffin JR, et al. A combined analysis of outcome following breast cancer: differences in survival based on BRCA1/BRCA2 mutation status and administration of adjuvant treatment. *Breast Cancer Res* 2004;6:R8–R17.
 49. Lips EH, Michaut M, Hoogstraat M, Mulder L, Besselink NJ, Koudijs MJ, et al. Next generation sequencing of triple negative breast cancer to find predictors for chemotherapy response. *Breast Cancer Res* 2015;17:134.
 50. Sharma P, Barlow W, Godwin AK, Knight L, Walker S, Kennedy R, et al. Impact of DNA repair deficiency signature on outcomes in triple negative breast cancer (TNBC) patients treated with AC chemotherapy (SWOG S9313) [abstract]. *J Clin Oncol* 2017;35:abstr 529.

Clinical Cancer Research

Pathological Response and Survival in Triple-Negative Breast Cancer Following Neoadjuvant Carboplatin plus Docetaxel

Priyanka Sharma, Sara López-Tarruella, José Angel García-Saenz, et al.

Clin Cancer Res 2018;24:5820-5829. Published OnlineFirst July 30, 2018.

Updated version Access the most recent version of this article at:
doi:[10.1158/1078-0432.CCR-18-0585](https://doi.org/10.1158/1078-0432.CCR-18-0585)

Supplementary Material Access the most recent supplemental material at:
<http://clincancerres.aacrjournals.org/content/suppl/2018/07/28/1078-0432.CCR-18-0585.DC1>

Cited articles This article cites 47 articles, 17 of which you can access for free at:
<http://clincancerres.aacrjournals.org/content/24/23/5820.full#ref-list-1>

Citing articles This article has been cited by 3 HighWire-hosted articles. Access the articles at:
<http://clincancerres.aacrjournals.org/content/24/23/5820.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://clincancerres.aacrjournals.org/content/24/23/5820>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.